

**PARTIAL MAYO SCORING INDEX
ASSESSMENT FOR ULCERATIVE COLITIS**

Date: / /
Patient name:
DOB:
UR number / address

Stool frequency (based on the past 3 days):

- | | POINTS | |
|--|--------|----------------------|
| <input type="checkbox"/> Normal number of stools | 0 | |
| <input type="checkbox"/> 1-2 stools/day more than normal | 1 | <input type="text"/> |
| <input type="checkbox"/> 3-4 stools / day more than normal | 2 | |
| <input type="checkbox"/> >4 stools / day more than normal | 3 | |

(normal = baseline number of stools e.g., before diagnosis of colitis)

**Rectal bleeding (based on the past 3 days
representing the most severe bleeding):**

- | | POINTS | |
|---|--------|----------------------|
| <input type="checkbox"/> None | 0 | |
| <input type="checkbox"/> Visible blood with stool less than half the time | 1 | <input type="text"/> |
| <input type="checkbox"/> Visible blood with stool half the time or more | 2 | |
| <input type="checkbox"/> Passing blood alone | 3 | |

(*note: a score of 3 requires patients to have at least 50% of bowel motions accompanied by visible blood and at least one bowel motion with blood alone)

Physician's global assessment

("based on patient's overall status on the day of the clinic visit")

- | | POINTS | |
|-----------------------------------|--------|----------------------|
| <input type="checkbox"/> Normal | 0 | |
| <input type="checkbox"/> Mild | 1 | <input type="text"/> |
| <input type="checkbox"/> Moderate | 2 | |
| <input type="checkbox"/> Severe | 3 | |

Total Partial Mayo Score (sum of above items)

- | | |
|---------------------------|----------------------|
| Remission = 0 to 1 | <input type="text"/> |
| Mild disease = 2 to 4 | |
| Moderate disease = 5 to 6 | |
| Severe disease = 7 to 9 | |